

A case report of severe panhypopituitarism in a newborn delivered by a women with Turner syndrome

Marta OLSZEWSKA¹, Grzegorz KIELBASA¹, Małgorzata WÓJCIK^{2,3},
Agata ZYGMUNT-GÓRSKA², Jerzy B. STARZYK^{2,3}

¹ Students Scientific Group, Department of Pediatric and Adolescent Endocrinology, Chair of Pediatrics, Institute of Pediatrics, Jagiellonian University Medical College, Krakow, Poland

² Children's University Hospital, Krakow, Poland

³ Department of Pediatric and Adolescent Endocrinology, Chair of Pediatrics, Institute of Pediatrics, Jagiellonian University Medical College, Krakow, Poland

Correspondence to: Małgorzata Wójcik MD., PhD.
Department of Pediatric and Adolescent Endocrinology, Chair of Pediatrics
Polish-American Pediatric Institute, Jagiellonian University Medical College,
Wielicka St. 265, 30-663 Krakow, Poland.
TEL: +48126581277; FAX: +486581005; E-MAIL: malgorzata.wojcik@uj.edu.pl

Submitted: 2015-12-23 Accepted: 2015-12-28 Published online: 2015-01-23

Key words: **Turner syndrome; pregnancy; pituitary stalk interruption syndrome; panhypopituitarism; case report**

Neuroendocrinol Lett 2015;36(8):734–736 PMID: 26921572 NEL360815C02 © 2015 Neuroendocrinology Letters • www.nel.edu

Abstract

Turner syndrome (TS) is a congenital disease caused by absence or structural abnormalities of sex chromosomes resulting in gonadal dysgenesis. Spontaneous pregnancies occur in 2–8% of patients, especially with mosaic karyotypes, however they are associated with increased risk of poor outcome both for mother and fetus. We report a 4-day-old male infant delivered by women with mosaic TS who was admitted to the pediatric intensive care unit and presented with severe panhypopituitarism as the early manifestation of pituitary stalk interruption syndrome (PSIS). To the best of our knowledge this is the first report of severe panhypopituitarism in a newborn borne by women with TS.

INTRODUCTION

Turner syndrome (TS) is a congenital disease caused by numerical or structural aberrations of sex chromosomes occurring in 1:2 000 to 1:2 500 liveborn females. It is usually characterized by short stature and gonadal dysgenesis resulting in delayed puberty, primary amenorrhea and infertility (Lacka 2005; Abir *et al.* 2001). Spontaneous puberty is observed in 15–30% of girls with TS and 2–20% of patients experience menarche without hormonal substitution. Spontaneous conceptions occur in 2–8% (Abir *et al.* 2001; El-Mansoury *et al.* 2007; Hagman *et al.* 2011; Birkebaek *et al.* 2002; Hewitt *et al.* 2013).

Pregnancies in TS women are associated with increased risk of poor outcome both for mother and fetus. Pregnancy-related complications in TS patients include preeclampsia, aortic dissection or rupture and spontaneous abortion. In children delivered by mothers with TS increased rate of prematurity, intrauterine growth restriction and chromosomal aberrations, especially TS and Down syndrome was observed (Abir *et al.* 2001; Hewitt *et al.* 2013). Congenital anomalies including ambiguous genitalia and hydrocephalus have been also reported (Birkebaek *et al.* 2002). To the best of our knowledge this is the first report of severe panhypopituitarism in a patient borne by women with TS.

CASE PRESENTATION

A 4-day-old male infant was admitted to the pediatric intensive care unit due to hypoglycaemia (2 mmol/L), myoclonic seizures and respiratory distress. He was delivered by women with TS diagnosed at the age of 36. She was referred for peripheral blood karyotype analysis due to primary infertility, but conceived spontaneously in the interterm before obtaining karyotype result, which revealed mosaic TS: 45,X/46,XX/47,XXX (92%/6%/2%). Her partner's karyotype was 46,XY.

The child was born at 36 weeks of gestation, via cesarean section due to decreased fetal heart rate variability in cardiotocography, with birth weight 2000 g and APGAR scores of 7 and 8 at 1 and 5 min, respectively. The patient's karyotype was 46,XY. Physical examination revealed jaundice and left-sided cryptorchidism. The child was diagnosed with early-onset sepsis (white blood cells: 20000/ μ L, C-reactive protein: 21.8 mg/L, interleukin 6: >1000 pg/mL), disseminated intravascular coagulation (platelet count: 102000/ μ L, APTT: 125.9 s, INR: 2.29, fibrinogen level: 1.15 g/L), prolonged jaundice with elevated hepatic enzymes (total bilirubin: 256 μ mol/L, direct bilirubin: 57.4 μ mol/L, aspartate transaminase: 205 IU/L, alanine transaminase: 132 IU/L) and diabetes insipidus (blood sodium level 147 mmol/L and serum osmolality 317 mOsm/kg). The child presented with persistent hypoglycaemia (min. 0.56 mmol/L). Hormonal analysis disclosed secondary hypothyroidism (TSH: 0.15 μ IU/mL, N: 0.8–9.1 μ IU/mL, fT4 6.4 pmol/L, N: 10–25 pmol/L), secondary adrenal insufficiency (ACTH <5 pg/mL, N: 10–60 pg/mL, blood cortisol:

2.9 ng/dL N: 50–230 ng/dL) and hypogonadotropic hypogonadism (FSH: 1.04 IU/L, N: 1.22–5.19 IU/L, LH: 2.42 IU/L, N: 4.85–10 IU/L, testosterone <0.1 ng/mL, N 0.2–0.49 ng/mL). Growth hormone deficiency was confirmed by test of nocturnal GH secretion and in stimulation test with arginine (max GH 6.32 ng/mL and 5.86 ng/mL respectively). Magnetic resonance imaging (MRI) of the brain revealed hypoplastic pituitary with thin pituitary stalk, lacking normal posterior pituitary signal and absence of septum pellucidum (Figures 1 and 2). The diagnosis of pituitary stalk interruption syndrome (PSIS) with early manifestation of severe panhypopituitarism was set. Replacement therapy with hydrocortisone, thyroxine, recombinant human growth hormone (rhGH) and desmopressin was implemented.

DISCUSSION

Spontaneous pregnancies are rarely observed in women with TS, especially over 34 years of age (Abir *et al.* 2001). Fertility without use of assisted reproductive technologies occur mostly in patients with mosaic karyotypes (Lacka 2005; Abir *et al.* 2001). This fact stays in line with the current evidence that only 46,XX germ cells in ovaries are able to complete meiosis (Modi *et al.* 2003). However, spontaneous pregnancies were reported also in simple X monosomic patients (El-Shawarby & Steer 2007; Mortensen *et al.* 2010). The most likely reason for that is, that the most frequently performed peripheral blood karyotype may not reflect chromosomal status of the ovarian tissue (Hewitt *et al.* 2013).

In most of the previously conducted studies pregnancies in TS women are regarded as high risk due




Fig. 1. Sagittal T1-weighted image showing hypoplastic anterior pituitary and absence of septum pellucidum.




Fig. 2. Coronal T1-weighted image – thin pituitary stalk and absence of septum pellucidum.

to baseline genetic load including thyroid dysfunction, congenital heart defects, higher susceptibility to obesity, diabetes mellitus and acquired cardiovascular diseases such as hypertension (Abir *et al.* 2001; Hewitt *et al.* 2013). Severe complications including fatal aortic dissection or rupture and heart failure have been described (Karnis *et al.* 2003). Nevertheless some of the studies do not confirm the increased maternal risk in these patients (Hagman *et al.* 2011).

Most of other authors emphasized, that pregnancies in TS patients are associated with increased rate of spontaneous abortion, which is likely caused by high prevalence of chromosomal aberrations in fetus, especially TS and Down syndrome (Abir *et al.* 2001; Hewitt *et al.* 2013). However, small uterus size and poor endometrial receptivity in TS women are another postulated reasons of miscarriages (Hewitt *et al.* 2013; Bakalov *et al.* 2007). The reported incidence of congenital anomalies in liveborn children of TS women ranges between 4.5% and 20% (Abir *et al.* 2001; Hagman *et al.* 2011, Hewitt *et al.* 2013).

To our knowledge it is the first report of PSIS manifested dramatically as a panhypopituitarism in a newborn delivered by women with TS. PSIS is defined as an association of hormonal deficiencies and radiological findings including thin or absent pituitary stalk, aplastic or hypoplastic anterior pituitary and ectopic or absent posterior pituitary. It can be associated with other mid-line defects, as in our patient presented with absence of septum pellucidum (Wang *et al.* 2014). PSIS is usually manifested as isolated growth hormone deficiency or multiple pituitary hormone deficiency with or without diabetes insipidus. The age of clinical manifestation ranges from neonates to adults. Our case report stays in line with other studies presenting persistent hypoglycaemia, prolonged jaundice, cryptorchidism and micropenis as typical signs of PSIS in neonates (Pinto *et al.* 1997). In older children short stature is regarded as the most common symptom (Ram *et al.* 2014).

Although in some cases of PSIS genetic etiology (mutations of HESX1, LHX4, OTX2, or SOX3) have been reported, in over 95% the mechanism of PSIS remains unknown (Reynaud *et al.* 2011; Yang *et al.* 2013). Considering the increased frequency of breech delivery and cesarean sections in children with PSIS, in most patients the role perinatal factors (trauma and hypoxia) such as in presented case is suggested (Wang *et al.* 2014).

Clinical manifestation of PSIS ranges in age of first symptoms and their severity with especially life-threatening course when occurring as panhypopituitarism in newborns. The genetic etiology of PSIS (mutations of HESX1, LHX4, OTX2, or SOX3) have been docu-

mented in less than 5% of cases. In most patients the mechanism of PSIS remains unknown and as in presented case requires further investigation. As maternal TS might be an essential risk factor for congenital abnormalities, infants borne by women with TS should remain under special medical supervision.

ACKNOWLEDGEMENTS

No funding to declare.

REFERENCES

- 1 Abir R, Fisch B, Nahum R, Orvieto R, Nitke S, Ben Rafael Z (2001). Turner's syndrome and fertility: current status and possible putative prospects. *Hum Reprod Update*. **7**: 603–610.
- 2 Bakalov VK, Shawker T, Cenicerio I, Bondy CA (2007). Uterine Development in Turner Syndrome. *J Pediatr*. **151**: 528–531.
- 3 Birkebaek NH, Crüger D, Hansen J, Nielsen J, Bruun-Petersen G (2002). Fertility and pregnancy outcome in Danish women with Turner syndrome. *Clin Genet*. **61**: 35–39.
- 4 El-Mansoury M, Barrenäs ML, Bryman I, Hanson C, Larsson C, Wilhelmsen L, *et al.* (2007). Chromosomal mosaicism mitigates stigmata and cardiovascular risk factors in Turner syndrome. *Clin Endocrinol (Oxf)*. **66**: 744–751.
- 5 El-Shawarby SA, Steer CV (2007). Spontaneous monozygotic monochorionic twin pregnancy in non-mosaic Turner's syndrome. *Int J Gynaecol Obstet*. **97**: 155.
- 6 Hagman A, Källén K, Barrenäs ML (2011). Obstetric outcomes in women with Turner karyotype. *J Clin Endocrinol Metab*. **96**: 3475–3482.
- 7 Hewitt JK, Jayasinghe Y, Amor DJ, Gillam LH, Warne GL, Grover S, *et al.* (2013). Fertility in Turner syndrome. *Clin Endocrinol (Oxf)*. **79**: 606–614.
- 8 Karnis MF, Zimon AE, Lalwani SI, Timmreck LS, Klipstein S, Reindollar RH (2003). Risk of death in pregnancy achieved through oocyte donation in patients with Turner syndrome: a national survey. *Fertil Steril*. **80**: 498–501.
- 9 Lacka K (2005). Turner's Syndrome – correlation between karyotype and phenotype. *Endokrynol Pol*. **56**: 986–993.
- 10 Modi DN, Sane S, Bhartiya D (2003). Accelerated germ cell apoptosis in sex chromosome aneuploid fetal human gonads. *Mol Hum Reprod*. **9**: 219–225.
- 11 Mortensen KH, Rohde MD, Uldbjerg N, Gravholt CH (2010). Repeated Spontaneous Pregnancies in 45,X Turner Syndrome. *Obstet Gynecol*. **115**: 446–449.
- 12 Pinto G, Netchine I, Sobrier ML, Brunelle F, Souberbielle JC, Brauner R (1997). Pituitary stalk interruption syndrome: a clinical-biological-genetic assessment of its pathogenesis. *J Clin Endocrinol Metab*. **82**: 3450–3454.
- 13 Ram N, Ali SA, Hussain SZ (2014). Pituitary stalk interruption syndrome presenting as short stature: a case report. *J Med Case Rep*. **8**: 445.
- 14 Reynaud R, Albarel F, Saveanu A (2011). Pituitary stalk interruption syndrome in 83 patients: novel HESX1 mutation and severe hormonal prognosis in malformative forms. *Eur J Endocrinol*. **164**: 457–465.
- 15 Wang Q, Hu Y, Li G, Sun X (2014). Pituitary stalk interruption syndrome in 59 children: the value of MRI in assessment of pituitary functions. *Eur J Pediatr*. **173**: 589–595.
- 16 Yang Y, Guo QH, Wang BA, Dou JT, Lv ZH, Ba JM, *et al.* (2013). Pituitary stalk interruption syndrome in 58 Chinese patients: clinical features and genetic analysis. *Clin Endocrinol (Oxf)*. **79**: 86–92.